

Comparative Studies on Physicochemical Properties and fatty Acids Composition of Seed Oil of *Jatropha curcas* and *Jatropha glauca*

Rania D.M. Albasha¹, Mawahib E.M. ElNour¹, Rajaa.S.M.Abadi ², Marmar A. El Siddig³

¹Department of Biology and Biotechnology, Faculty of Science and Technology, AL Neelain University.

²Department of Chemistry, Faculty of Science and Technology, AL Neelain University

³Department of Botany, Faculty of Science, Khartoum University

Raniaelbasha90@gmail.com

aelnourmawahib@gmail.com

RajaaSatti@hotmail.com

marmarelsiddig@hotmail.com

Abstract— (The aim of this study was to investigate the physical and chemical properties of the seeds oil of *Jatropha curcas* (L) and *Jatropha glauca* collected from two different regions of Sudan. The physicochemical analyses were carried out using standard analytical methods. Results showed that the specific gravity and refractive index @ 250C were 0.943 and 1.473 for *J.curcas* compared to 0.951 and 1.470 for *J.glauca*, respectively. The recorded acid values, saponification number, peroxide values and % of free fatty acids (mgKOHg-1 oil) were, in order, 9.537 and 14.025; 193.40 and 206.70; 1.93 and 2.3; 4.769 and 7.012 for *J.curcas* and *J.glauca*, respectively. On the other hand, iodine values (g100g-1oil) were 114.64 and 117.08 for *J.curcas* and *J.glauca*, respectively. Fatty acids composition of the two species was revealed using the gas chromatography-mass spectrometry (GC-MS) technique. The dominant unsaturated fatty acids of *Jatropha curcas* seeds oil were oleic acid (37.03%) and linoleic acid (21.60%), while the saturated fatty acids were stearic acid (10.95%), palmitic acid (17.81%) and archidic (1.00%). Comparatively, the unsaturated fatty acids of *J. glauca* were linoleic acid (51.74%) and oleic acid (15.37%) whereas the saturated fatty acids of this species were stearic acid (13.18%), palmitic acid (14.92%) and archidic(1.14%). The results of chemical and physical properties of seed oil of the two *jatropha* species suggest that the seed oil may be suitable for industrial purposes such as production of soaps, paints and lubricants. At the same time, results obtained from (GC) analysis revealed that the types of fatty acids of *J. curcas* seed oil may be more suitable for production of biodiesel than that of other species.

Index Terms— *Jatropha curcas*, *Jatropha glauca*, Physicochemical Properties, Seed Oil , Fatty Acids.

I. INTRODUCTION

The genus *Jatropha* has 175 known species of the plant belonging to the family Euphorbiaceae. These species are mainly grown in warm temperate regions and seasonally dry tropics. Africa counts 70

native species and Madagascar has 1 endemic. Several other *Jatropha* species occur in the same region as *Jatropha glauca* and have medicinal uses [1],[2]mentioned that there are a number of non edible, tree based oil seeds available in many countries around the world from which biodiesel can be produced.

Jatropha curcas L. and *Jatropha glauca* are multipurpose shrubs with a variety of applications and enormous economic potentials for their seed oil, which can be converted into biodiesel an alternative to petro-diesel. As the future lack of petroleum is a current concern, bio-diesel seems to be “part of the solution”, by replacing partial or totally petro-diesel fuel in diesel engines. This reason, added to an increasing environmental concern, creates a scenario in which biodiesel production is expected to have a big development over the next few years [3].

Jatropha glauca occurs in Sudan, Eritrea, Ethiopia, Djibouti and Somalia, and extends to Yemen and Saudi Arabia. Botanically it is describe as small monoecious shrub up to 1 m tall with smooth, pale branches; stems and leaves glabrous to shortly hairy .*Jatropha glauca* occurs in open bush land, extending to semi-desert conditions, on lava and limestone, from sea-level up to 1000 m altitude. *Jatropha glauca* is conceder as natural vegetation, dominant plant to western plain of the Red Sea

Hills. Sandy soil is the dominant soil type in the coastal and western plains. Sandy soils are characterized by their low fertility and high infiltration rate, hence unsuitable for cultivation. Natural vegetation is sparse and the grazing capacity is relatively low [4]. The region experiences both winter and summer rains. It is estimated that 98% of the region receive less than 200 mm. of annual rainfall. An average annual temperature of western and southern plain is above 22°C [5]. *Jatropha glauca* is relatively common in its distribution area and is not browsed by livestock, it is therefore not likely to be threatened by genetic erosion [6].

The centre of origin of *J. curcas* is still debatable, but it is believed to be native of Mexico and Central American region. It has been introduced to Africa and Asia in the new world. It is cultivated in many parts of the tropics and subtropics as a hedge crop and for traditional use [7]; [8]. *Jatropha* occurs mainly at lower altitudes (0-500 m) in areas with average annual temperatures well above 20°C but can grow at higher altitudes and tolerates slight frost. It grows on well-drained soils with good aeration and is well adapted to marginal soils with low nutrient content. The current distribution shows that introduction has been most successful in the drier regions of the tropics and can grow under a wide range of rainfall regimes [9].

The literature indicates that the fatty acid methyl ester of *Jatropha curcas* is one of the 26 fatty acid methyl esters of oil that are most suitable for biodiesel product [10].

[11] Revealed that the fatty acid composition of *J. curcas* oil was analyzed by gas chromatography, major long chain fatty acids present in the *J. curcas* oil which are palmitic acid (16.69%), stearic acid (7.67%), oleic acid (40.39%) linoleic acid (33.09%) and Linolenic acid (0.28%). *J. curcas* oil contains high percentage of unsaturated fatty acid, which is about 75.64%. Also [12] revealed that *Jatropha curcas* (L) oil is rich in unsaturated fatty acids especially oleic acid (52.27%) and linoleic acid (27.87%). The dominant saturated fatty acids were palmitic acid (14.24%) and stearic acid (5.15%). These results suggest that *Jatropha curcas* (L) seed oil may not be suitable for human

consumption except it is subjected to detoxification and purification before use, but may be suitable for industrial purposes such as production of soaps, paints and lubricants. The physiochemical properties, energy value, fatty acid composition and oil content of the seeds of *Jatropha curcas* were investigated by [13]; [14]; [15]. These physical and chemical properties of *Jatropha curcas* oil are strongly influenced by the processing, season, climate and geography during the growth of the seed, storage etc. The rainfall, soil type, nutrition content of the soil, temperature and other factors vary from region to region and hence the physical and chemical properties of the *Jatropha curcas* oil produced are region specific [16]. Research of seed oil of *Jatropha curcas* and *Jatropha glauca* is not available in Sudan. Therefore, it is of great interest to evaluate the physical and chemical properties of the seed oil of these promising plants.

MATERIAL AND METHODS

A. 2.1 Plant materials:

Healthy seeds of *Jatropha curcas* were collected from *Jatropha curcas* plants cultivated in Alrwakeep research station, National Center for Research, Khartoum, Sudan, whereas the healthy seeds of *Jatropha glauca* were collected from *Jatropha glauca* plant grown naturally in Sinkat (valleys), Red Sea State, Sudan. The seeds for both species were identified at the Department of Biology & Biotechnology, Faculty of Science & Technology, AlNeedain University.

B. 2.2 Soxhlet extraction:

The method described by [17] with slight modification was used. The clean dry seeds (25 g) of *Jatropha curcas* and *Jatropha glauca* were grounded using mortar and pestle then defatted in a soxhlet apparatus. The extraction carried out by using petroleum ether (60-80°C). The process continued for 3 h. Solvent removed by vacuum evaporation in room temperature. Each extraction was run in triplicate and yield % was determined.

C. 2.3 Gas Chromatography-Mass Spectrometry (GC-MS) condition for analysis of fatty acid:

Methyl esters were prepared from total lipids by the method of [18]. These fatty acids methyl esters

were analyzed by gas chromatography, Shimadzu series (GC.MS.QP.2010), equipped with mass spectrometer detector and Rtx.50 column having internal diameter 0.25mm and length 30cm. Fatty acids profile obtained through this gas chromatography with relevant standards.

D. 2.4 Physicochemical analysis:

The specific gravity of the *J. curcas* and *J. glauca* oil were determined with specific gravity bottle method as described by [19]. The refractive indices of the oils were read at 250C using an Abbe Refractometer [20].

Iodine value determined by Hanus iodine method [21], while saponification values, acid values, and peroxide values were determined according to [18]. The amount of free fatty acid (FFA) was calculated as being equivalent to half the value of acid value. All the analyses were done in triplicate and reagents used were of analytical.

RESULTS AND DISCUSSION:

Data in Table 1 represented the physicochemical properties of *J. curcas* and *J. glauca* seed oil. The oil extracted from the two species of *Jatropha* using petroleum ether was transparent liquid at room temperature. The oil extracted from *J. curcas* seeds has bitter taste which may be due to the present of alkaloid; it has agreeable odour and light yellowish colour. The extracted oil of *J. glauca* has also bitter taste, pleasant odour and it was golden yellow in colour. The percentage oil yield of *J. curcas* and *J. glauca* were 28 and 26%, respectively. These results are contrary to that reported by [22], where the seed oil yield of *J. curcas* oil was 52.75%. Specific gravity *J. curcas* seed oil at 250C was 0.94 while it was 0.95 for *J. glauca*; these values showed consistency with studies of [23] and [22]. The refractive indices at 250C were 1.473 and 1.470 for *J. curcas* and *J. glauca*, respectively. These values were in agreement with that reported by [24]. The physical properties of the oil extracted from the two species were approximately similar (Table 1). On the other hand the chemical properties of the oil of two species were different as shown in the table 1. An acid value of the seed oil of *J. curcas* was 9.53 mg KOH/g-1 oil, this value showed

inconsistency with that obtained by [22] and [24], where their findings range between 36.20-36.46 mg KOH/g. This may be due to the difference in the climatic conditions [25]. The acid value of *J. glauca* oil showed higher value than that of *J. curcas* (14.02 mg KOH/g-1 oil). The free fatty acids (FFAs) of *J. curcas* oil were 4.77 mg KOH/g-1 oil. This result is quite consistent with the values obtained by previous researchers [12]. On the other hand, the FFAs of *J. glauca* oil were 7.01 mg KOH/g-1 oil. High FFAs (%wt) cause soap formation during alcoholysis process and lead to difficulties in separation of biodiesel from its byproducts; as a result it reduced the biodiesel yield [26]. Peroxide value were 1.93 and 2.30 mg KOH/g-1 oil and iodine value were 114.64 and 117.08 g/100g-1 oil for *J. curcas* and *J. glauca* oil, respectively (Table 1). Our results agree with that obtained by [12]. A low peroxide value, as seen in the present study, increases the suitability of the oil for a long-time storage due to a low level of oxidative and lipolytic activities. Iodine value is the measure of degree of unsaturation of the oil. Higher iodine value indicates higher unsaturation of fats and oils [27]. Saponification number of *J. curcas* oil recorded in this research was 193.40 mg KOH/g-1 oil. This high value was consistent with the findings in the previous literatures [17]; [22], who reported that *Jatropha curcas* oil is usually associated with high saponification value. Such high value establishes the fact that *Jatropha curcas* oil contains normal triglycerides. In this study, *J. glauca* seed oil gave higher amount of saponification number (206.7 mg KOH/g-1 oil) than that of *Jatropha curcas* seed oil. To the best of our knowledge, there is no documented study on seed oil extracted from *J. glauca*. Figures (1 and 2) as well as Table 2 represent the composition of fatty acids of seed oil of *J. curcas* and *J. glauca*, respectively, as determined by the gas chromatography-mass spectrometry. The properties of the triglyceride and the biodiesel fuel are determined by the amount of each fatty acid that is present in the molecule. Chain length and the number of double bonds determine the physical characteristics of both fatty acids and triglyceride [17]. There are three main types of fatty acids present in a triglyceride: Saturated

(CN:0), monounsaturated (CN:1) and polyunsaturated (CN:2,3) and an ideal vegetable oil as a potential feedstock for biodiesel production should have low saturated and polyunsaturated fatty acid and be high in monounsaturated fatty acid [28]. The results of this study revealed that the *Jatropha curcas* consist of saturated fatty acid especially palmitic acid (17.81%), stearic acid (10.95%) and arachidic acid (1.00%) whereas monounsaturated fatty acid was oleic acid (37.03%) and polyunsaturated fatty acid was linoleic acid (21.60%). This finding of monounsaturated fatty acid of seed oil *Jatropha curcas* was in accordance with the finding of [22] and approximately agree with that obtained by [11], but the amount of saturated fatty acid and polyunsaturated fatty acid were lower than that obtained by these researches. According to [28] the *Jatropha curcas* cultivated in Sudan, Khartoum, may be more suitable for biodiesel production. The prevalence of the unsaturated fatty acids and high values of the iodine index indicate that the *J. curcas* oil is of the unsaturated type [27].

The saturated fatty acids of *Jatropha glauca* were palmitic acid (14.92%), stearic acid (13.18%) and arachidic acid (1.14%). The results of saturated fatty acids recorded in the present study for *J. glauca* were approximately within the range of that of *Jatropha curcas*. Monounsaturated fatty acid of *J. glauca* was oleic acid (15.37%) and polyunsaturated was linoleic acid (51.74%). *J. glauca* oil contains high percentage of polyunsaturated fatty acid. Oil containing high amount of polyunsaturated fatty acids tend to exhibit poor oxidation stability, and may not be useful at low temperatures due to a high pour points, but can find an application in the surface coating industries [29]; [30]; and [24].

CONCLUSION:

J. curcas seed oil contained high percentage of monounsaturated fatty acid which is about 37.03%. On the other hand *J. glauca* showed high percentage of polyunsaturated fatty acid (51.74%). The results of this investigation suggest that *J. curcas* and *J. glauca* seed oil have great potentials for future industrial oil seeds crops.

REFERENCES

- [1] Dehgan, B. (1984). Phylogenetic significance of interspecific hybridization in *Jatropha* (Euphorbiaceae). *Systematic Botany*. 9 :467-478.
- [2] Kayashri, P. KJeyashoke, N. Krisangkura, K. (1996). Survey of seed oils for use as diesel fuels. *J. Am. Oil Chem. Soc.* 73 (4) 471-474.
- [3] Montes, J.M., Rodríguez Aliciardi, M., Vaca Chávez J., Guzmán, C. and Calandri, E. (2011). Characterization of *Jatropha curcas* L. Seed and its Oil, from Argentina and Paraguay. *The Journal of the Argentine Chemical Society*. 98:1-9.
- [4] Barbour, K.M. (1961). *The Republic of the Sudan: regional geography*. University of London Press, London.
- [5] ElTom, M.A. (1991). *The Climate of the Red Sea Region of the Sudan*. Proceedings of the (RESAP) workshop, Khartoum, Sudan.
- [6] Schmelzer G.H., Gurib-Fakim A., Arroo, R., Bosch, C.H., de Ruijter, A., Simmonds MSJ, eds. (2008) *Plant Resources of Tropical Africa 11(1) – Medicinal Plants 1*. Backhuys Publishers, Wageningen, Netherlands.
- [7] Heller, J. (1996). Promoting the conservation and use of under utilized and neglected crops. 1. Physic nut *Jatropha curcas* L. *International Plant Genetic Resources Institute*, Rome. 66pp.
- [8] Kumar, A. and Sharma, S. (2008). An evaluation of multipurpose oil seed crop for industrial uses: A Review. *Industrial Crops and Products* 28 :1-10.
- [9] Katwal R.P.S. and. Soni, P.L (2003). Biofuels: an opportunity for socioeconomic development and cleaner environment. *Indian Forester* 129: 939–949.
- [10] Mohibbe, A.M. Waris, A. and Nahar, N.M. (2005). Prospects and potential of fatty acid methyl esters of some non-traditional seed oils for use of biodiesel in India, *Biomass Bioenergy* 29 :293-302.
- [11] Archana Joshi, Pankaj S. and Bachheti, R.K. (2011). Physicochemical characterization of seed oil of *Jatropha curcas*. *International journal of applied biology and pharmaceutical technology*. 2: (2) 0976-4550.
- [12] Ugbogu, A. E., Akubugwo, E. I., Uhegbu, F. O., Chinyere, C. G., Ugbogu, O. C. and Oduse, K. A. (2014). Quality assessment profile of *Jatropha curcas* (L) seed oil from Nigeria. *International Food Research Journal*. 21(2): 735-741.
- [13] Pramanik, K. (2003). Properties and use of *Jatropha curcas* oil and diesel fuel blends in compression ignition engine, *Renewable Energy* 28: 239-248.
- [14] Shah, S. Sharma, A. Gupta, M.N. (2004). Extraction of oil from *Jatropha curcas* L. seed kernels by enzyme assisted three phase partitioning, *Industrial crops and products* 20: 275-279.
- [15] Tiwari, A.K. Kumar, A. Raheman, H. (2007). Biodiesel production from *Jatropha* oil (*Jatropha curcas*) with

high free fatty acids: An optimized process, Biomass Bioenergy 31(8): 569-575.

[16] Bojan, S.G and Durairaj, S.K.(2012). Producing Biodiesel from High Free Fatty Acid *Jatropha curcas* Oil by a Two Step Method- an Indian Case Study. Journal of Sustainable Energy & Environment 3 : 63-66.

[17] Akbar, E. Yaakob, Z. Kamarudin, S. Ismail, M. (2009). Characteristics and Composition of *Jatropha curcas* oil seed from Malaysia and its potential as Biodiesel Feedstock. Eur. J. scientific Res. 29: 396-403.

[18] AOAC, 1990. Association of Official Analytical Chemist. 14th Edn., AOAC Arlington. USA.

[19] Pearson, D. 1976. The Chemical Analysis of Food. Churchill, Livingstone pp: 488-496.

[20] Singhal, S.C. & Sekiya J. 2003. Modern Technology in the Oils and Fats Industry. 2nd (ed.) New Delhi: AOSC-OTA13.

[21] Association of Official Agricultural Chemists (AOAC) (1975). Official Methods of Analysis. Washington, 12th edition, pp. 488-492.

[22] Umaru, M. and Aberuagba, F.(2012). Characteristics of a Typical Nigerian *Jatropha curcas* oil Seeds for Biodiesel Production. Research Journal of Chemical Sciences 2 (10): 7-12.

[23] Belewu M.A, Adekola F.A., Adebayo G.B., Ameen O.M., Muhammad N.O., Olaniyan A.M., Adekola O.F. and Musa A.K., (2010). Physico-Chemical characteristic of oil and biodiesel from Nigerian and Indian, *Jatropha curcas* seeds, Int. J of Bio and Chem. Sci., 4(2), 524-529.

[24] Nayak, B.S. and Patel, K.N. (2010). Physicochemical Characterization of Seed and Seed Oil of *Jatropha curcas* L. collected from Bardoli (South Gujarat). Sains Malaysia 39: 951-955.

[25] Najda, A., Almehehdi, A.F. and Zabar, A.F. (2013). Chemical composition and nutritional value of *Jatropha curcas* L. leaves. Journal of Genetic and Environmental Resources Conservation 1(3):221-226.

[26] Freedman B., Pyrd E.H., Mounts T.L., (1984). Variables affecting yields of fatty esters from Transesterified vegetable oils, JA OCS 61,1638-1643.

[27] Nzikou, J.M., Matos, L., Moussounga, J.E., Ndangui, C.B., Kimbonguila, A., Silou, T.H., Linder, M. and Desobry, S. (2009). Characterization of *Moringa oleifera* seed oil variety congo-brazzaville. Journal of Food Technology 7: 59-65.

[28] Gunstone, F.D.,(2004). Rapeseed and Canola Oil: Production, Processing, properties and uses London: Black well Publishing Ltd .

[29] Augustus, G.D.P.S., Jayabalan M. and Seiler, G.J. (2002). Evaluation and bio-induction of energy components of *Jatropha curcas*. Biomass and Bioenergy 23: 161-164.

[30] Emil, A., Yaakob, Z., Kumar, M.N.S., Jahim, J.M. and Salimon, J. (2010). Comparative evaluation of physicochemical properties of *Jatropha* seed oil from Malaysia,

Indonesia and Thailand. Journal of the American Oil Chemists' Society 87:689-696]

I. Table 1: Physical and chemical properties of seed oil of *J. curcas* and *J. glauca*

Physicochemical properties		<i>J. glauca</i>	<i>J. curcas</i>
Physical properties			
Organoleptic	Yield	26%	28.8%
	Taste	Bitter	Bitter
	Odour	Pleasant	Agreeable
	Colour	Golden yellow	Light yellowish
Appearance at room temperature		transparent liquid	transparent liquid
Specific gravity at 250C		0.951	0.943
Refractive index at 250C		1.470	1.473
Chemical properties			
Acid value		14.025	9.537
Iodine value		117.08	114.64
Saponification value		206.7	193.5
% FFA		7.0125	4.769
Peroxide value		2.3	1.93

Table 2: Fatty acids composition of seed oil of *J. curcas* and *J. glauca*

No.	Name of The Compound	Composition (%)	
		<i>J. curcas</i>	<i>J. glauca</i>
1	Palmitic acid	17.81	14.92
2	Stearic acid	10.95	13.18
3	Arachidic acid	1.00	1.14
4	Oleic acid	37.03	15.37
5	Linoleic acid	21.60	51.74

