

MAXIMAL AND MINIMAL IDEALS IN TRANSFORMATION SEMIGROUPS WITH INVARIANT SETS

Attapol Praleah, Jintana Sanwong

Mathematics Department
Faculty of Science, Chiang Mai University
Chiang Mai, Thailand
attapol_praleah@outlook.com

Abstract—Let X be a set and $T(X)$ denote the semigroup (under composition) of transformations from X into itself. For a fixed nonempty subset Y of X , let

$$S(X, Y) = \{\alpha \in T(X) : Y\alpha \subseteq Y\}.$$

Then $S(X, Y)$ is a semigroup of total transformations of X which leave a subset Y of X invariant. In this paper, existence and uniqueness of maximal and minimal ideals of $S(X, Y)$ are proved. Moreover, we present a maximal congruence on $S(X, Y)$ when X is a finite set.

Keywords—maximal ideals, minimal ideals, transformation semigroups with invariant sets.

I. INTRODUCTION

Let X be a set and $\emptyset \neq Y \subseteq X$. The semigroup we consider is $S(X, Y)$ consists of all mappings in $T(X)$ which leave $Y \subseteq X$ invariant. K. D. Magill [5] introduced and studied the semigroup $S(X, Y)$ in 1996. In fact, if $Y = X$, then $S(X, Y) = T(X)$. So we may regard $S(X, Y)$ as a generalization of $T(X)$. In 2005, S. Nenthein, P. Youngkhong, and Y. Kemprasit [7] showed that $S(X, Y)$ is a regular semigroup if and only if $X = Y$ or Y contains exactly one element, and $\text{Reg } S(X, Y) = \{\alpha \in S(X, Y) : X\alpha \cap Y = Y\alpha\}$ is the set of all regular elements of $S(X, Y)$. Moreover, they counted the numbers of regular elements in $S(X, Y)$ for a finite set X . The numbers were given in terms of the cardinalities of X and Y . Later in 2013, W. Choomanee, P. Honyam and J. Sanwong [1] studied left regular, right regular and intra-regular elements of $S(X, Y)$ and consider the relationships between these elements. Moreover, they counted the number of left regular elements of $S(X, Y)$ when X is a finite set.

As far back in 1952, Malcev [6] determined ideals of $T(X)$. In 2011 P. Honyam and J. Sanwong [4] characterized when $S(X, Y)$ is isomorphic to $T(Z)$ for some set Z and prove that every semigroup A can be embedded in $S(A^1, A)$. Then they described Green's relations and ideals on $S(X, Y)$ and applied these results to obtain its group H -classes and ideals.

In this paper, we determine maximal and minimal ideals of $S(X, Y)$. We also present a maximal congruence on $S(X, Y)$ when X is a finite set.

II. PRELIMINARIES AND NOTATIONS

In this section, we list some known results, definitions and notations that will be used throughout this paper.

Let X be a set and Y a nonempty subset of X . Then $S(X, Y)$ is a semigroup with identity 1_X , the identity map on X . Green's relation on $S(X, Y)$ are given by P. Honyam and J. Sanwong [4], which are needed in characterizing ideals on $S(X, Y)$.

Lemma 2.1. [4] Let $\alpha, \beta \in S(X, Y)$. Then

(1) $\alpha L \beta$ if and only if $X\alpha = X\beta$ and $Y\alpha = Y\beta$;

(2) $\alpha R \beta$ if and only if $\pi_\alpha = \pi_\beta$ and $\pi_\alpha(Y) = \pi_\beta(Y)$

(3) $\alpha J \beta$ if and only if $|X\alpha| = |X\beta|, |Y\alpha| = |Y\beta|$ and

$$|X\alpha \setminus Y| = |X\beta \setminus Y|$$

Let p be any cardinal number such that

$$p' = \min\{q : q > p\}.$$

Note that p' always exists since the cardinals are well-ordered and when p is finite we have $p' = p + 1 =$ the successor of p .

To describe ideals of $S(X, Y)$, we let $|X| = a$, $|Y| = b$ and $|X \setminus Y| = c$. For each cardinals r, s, t such that $2 \leq r \leq a'$, $2 \leq a \leq b'$ and $1 \leq t \leq c'$, we define

$$S(r, s, t) = \{\alpha \in S(X, Y) : |X\alpha| < r, |Y\alpha| < s \text{ and } |X\alpha \setminus Y| < t\}.$$

Theorem 2.2. [4] The set $S(r, s, t)$ is an ideal of $S(X, Y)$.

To obtain ideals of $S(X, Y)$, we need the following notation. Let Z be a nonempty subset of $S(X, Y)$, and let

$$K(Z) = \{\alpha \in S(X, Y) : |X\alpha| \leq |X\beta|, |Y\alpha| \leq |Y\beta| \text{ and}$$

$$|X\alpha \setminus Y| \leq |X\beta \setminus Y| \text{ for some } \beta \in Z\}.$$

Then we see that $Z \subseteq K(Z)$ and $Z_1 \subseteq Z_2$ implies that $K(Z_1) \subseteq K(Z_2)$.

Theorem 2.3. [4] The ideals of $S(X, Y)$ are precisely the set $K(Z)$ for some nonempty subset Z of $S(X, Y)$.

Let $G(A)$ be the group of permutation on the set A . Define

$$G(X, Y) = \{\alpha \in G(X) : \alpha|_Y \in G(Y)\},$$

Where $Y \subseteq X$ and $\alpha|_Y$ is the restriction of α on the set Y .

Then $G(X, Y)$ is a subgroup of the permutation group $G(A)$.

If X is a finite set with n elements and Y a nonempty subset of X with m elements, then we define

$$J_{r,s,t} = \{\alpha \in S(X,Y) : |X\alpha| = r, |Y\alpha| = s \text{ and } |X\alpha \setminus Y| = t\}$$

and

$$J_k = \{\alpha \in S(X,Y) : |X\alpha| = k\}$$

where $1 \leq r \leq n$, $1 \leq s \leq m$, $0 \leq t \leq n-m$ and $1 \leq k \leq n$. Thus $J_{r,s,t}$ is a J -class of $S(X,Y)$, J_1 is the set of all constant maps with image in Y and $J_n = G(X,Y)$.

The following convenient notation will be used in this paper: given $\alpha \in S(X,Y)$ we write

$$\alpha = \begin{pmatrix} X_i \\ a_i \end{pmatrix},$$

and take as understood that the subscript i belongs to some (unmentioned) index set I , the abbreviation $\{a_i\}$ denote $\{a_i : i \in I\}$, and that $X\alpha = \{a_i\}$ and $a_i\alpha^{-1} = X_i$.

With the above notation, for any $\alpha \in S(X,Y)$ we can write

$$\alpha = \begin{pmatrix} A_i & B_j & C_k \\ a_i & b_j & c_k \end{pmatrix},$$

where $A_i \cap Y \neq \emptyset$; $B_j, C_k \subseteq X \setminus Y$; $\{a_i\} \subseteq Y$; $\{b_j\} \subseteq Y \setminus \{a_i\}$

and $\{c_k\} \subseteq X \setminus Y$. Here, I is a nonempty set, but J or K can be empty.

III. RESULTS

From now on, we let X be a finite set with n elements and Y a nonempty subset of X with m elements. Let \mathcal{I} be the set of all ideals of $S(X,Y)$. Then (\mathcal{I}, \subseteq) is a partially ordered set with the following property.

Theorem 3.1. J_1 is a minimum ideal of $S(X,Y)$.

Proof: We prove that $J_1 = S(2,2,1)$. It is clear that $J_1 \subseteq S(2,2,1)$. Let $\alpha \in S(2,2,1)$. Then $|X\alpha| < 2, |Y\alpha| < 2, |X\alpha \setminus Y| < 1$ and thus $|X\alpha| = 1 = |Y\alpha|$. So α is a constant map and $\alpha \in J_1$. Therefore, J_1 is an ideal of $S(X,Y)$. To show that J_1 is a minimum ideal, let I be an ideal of $S(X,Y)$ and $\beta \in J_1$. Then there exists $\emptyset \neq Z \subseteq S(X,Y)$ such that $I = K(Z)$. Let $\gamma \in Z$. Then $|X\gamma|, |Y\gamma| \geq 1$, so $|X\beta| = 1 \leq |X\gamma|, |Y\beta| = 1 \leq |Y\gamma|$ and $|X\beta \setminus Y| = 0 \leq |X\gamma \setminus Y|$. Thus $\beta \in I$, i.e., $J_1 \subseteq I$ as required. +

Lemma 3.2. If $|Y| = 1$ and $J_{2,s,t} \neq \emptyset$, then $J_1 \cup J_{2,s,t}$ is an ideal of $S(X,Y)$ if and only if $s = 1 = t$.

Proof: Assume that $|Y| = 1$ and $J_{2,s,t} \neq \emptyset$. Suppose that $J_1 \cup J_{2,s,t}$ is an ideal of $S(X,Y)$. Let $\alpha \in J_{2,s,t}$. Since $Y\alpha \subseteq Y$, we have $1 \leq |Y\alpha| \leq |Y| = 1$, so $|Y\alpha| = 1$ which implies that $s = |Y\alpha| = 1$. Since $|X\alpha| = 2$ and $Y = Y\alpha$, we have $|X\alpha \setminus Y| = |X\alpha \setminus Y\alpha| = 2 - 1 = 1$, that is $t = 1$.

Conversely, assume that $s = 1 = t$. First we show that $J_1 \cup J_{2,1,1} = S(3,2,2)$. Let $\alpha \in J_1 \cup J_{2,1,1}$. Then $\alpha \in J_1$ or $\alpha \in J_{2,1,1}$. If $\alpha \in J_1$, then $|X\alpha| = 1 < 3, |Y\alpha| \leq |X\alpha| = 1 < 2$ and $|X\alpha \setminus Y| = 1 - 1 = 0 < 2$. thus $\alpha \in S(3,2,2)$. If $\alpha \in J_{2,1,1}$,

then $|X\alpha| = 2 < 3, |Y\alpha| = 1 < 2$ and $|X\alpha \setminus Y| = 1 < 2$. Thus $\alpha \in S(3,2,2)$. For the other containment, let $\alpha \in S(3,2,2)$. Then $|X\alpha| \leq 2, |Y\alpha| \leq 1$ and $|X\alpha \setminus Y| \leq 1$. If $|X\alpha| = 1$, then $\alpha \in J_1$. if $|X\alpha| = 2$, then $|X\alpha \setminus Y| = |X\alpha \setminus Y\alpha| = 2 - 1 = 1$. Then $\alpha \in J_{2,1,1}$. Hence $J_1 \cup J_{2,1,1} = S(3,2,2)$ is an ideal. +

Since J_1 is the minimum ideal of $S(X,Y)$, we define a minimum ideal in $S(X,Y)$ as follows. An ideal $J_1 \emptyset I$ of $S(X,Y)$ is a minimal ideal if J is an ideal such that $J_1 \emptyset j \subseteq I$, then $J = I$.

Theorem 3.3. If $|Y| = 1$, then $J_1 \cup J_{2,1,1}$ is the unique minimal ideal of $S(X,Y)$.

Proof: Suppose that $Y = \{a\}$. By Lemma 3.2, we have $J_1 \cup J_{2,1,1}$ is an ideal of $S(X,Y)$. Next, we show that $J_1 \cup J_{2,1,1}$ is a minimal ideal of $S(X,Y)$. Let J be an ideal of $S(X,Y)$ such that $J_1 \subseteq J \subseteq J_1 \cup J_{2,1,1}$. Suppose that $J \emptyset J_1 \cup J_{2,1,1}$. It is clear that $J_1 \subseteq J$. By assumption, we have exists $\alpha \in J_{2,1,1}$ but $\alpha \notin J$. We show that $J \subseteq J_1$ by supposing this is false, so $J \not\subseteq J_1$. Then there exists $\beta \in J$, but $\beta \notin J_1$. Since $\alpha, \beta \in J_{2,1,1}$, we can write

$$\alpha = \begin{pmatrix} A & X \setminus A \\ a & b \end{pmatrix}$$

Where $Y \subseteq A, a \in Y, b \in X \setminus Y$ and

$$\beta = \begin{pmatrix} B & X \setminus B \\ a & c \end{pmatrix}$$

Where $Y \subseteq B, c \in X \setminus Y$. Let $\gamma, \theta \in S(X,Y)$ be defined by

$$\gamma = \begin{pmatrix} A & X \setminus A \\ u & v \end{pmatrix}, \quad \theta = \begin{pmatrix} Y & X \setminus Y \\ a & b \end{pmatrix},$$

Where $u \in B \cap Y, v \in X \setminus B$. Consider

$$\begin{aligned} \gamma\beta\theta &= \begin{pmatrix} A & X \setminus A \\ u & v \end{pmatrix} \begin{pmatrix} B & X \setminus B \\ a & c \end{pmatrix} \begin{pmatrix} Y & X \setminus Y \\ a & b \end{pmatrix} \\ &= \begin{pmatrix} A & X \setminus A \\ a & b \end{pmatrix} = \alpha. \end{aligned}$$

Then $\alpha = \gamma\beta\theta \in J$, which is a contradiction. So $J = J_1$.

Hence, $J_1 \cup J_{2,1,1}$ is a minimal ideal of $S(X,Y)$. Finally, we show that $J_1 \cup J_{2,1,1}$ is a unique minimal ideal of $S(X,Y)$. We show that $M = N$. Since N is an ideal of $S(X,Y)$, we get that $N = K(Z)$ for some $\emptyset \neq Z \subseteq S(X,Y)$. Since $J_1 \emptyset N$, there exists $\alpha \in N$ with $|X\alpha| \geq 2$. Since $\alpha \in N = K(Z)$, we obtain that $|X\alpha| \leq |X\beta|, |Y\alpha| \leq |Y\beta|$ and $|X\alpha \setminus Y| \leq |X\beta \setminus Y|$ for some $\beta \in Z$. Let $\gamma \in J_{2,1,1}$. Then $|X\gamma| = 2$ and so $|X\gamma| \leq |X\alpha| \leq |X\beta|$. Since $|Y| = 1$, we have $|Y\gamma| = 1 = |Y\alpha| \leq |Y\beta|$ and $|X\gamma \setminus Y| = 1 \leq |X\alpha \setminus Y| \leq |X\beta \setminus Y|$. Then $\gamma \in K(Z) = N$. Thus $J_{2,1,1} \subseteq N$ and so $J_1 \cup J_{2,1,1} \subseteq N$ which implies that $M = N$.

Lemma 3.4. If $|Y| > 1$ and $J_{2,s,t} \neq \emptyset$, then $J_1 \cup J_{2,s,t}$ is an ideal of $S(X,Y)$ if and only if $s = 1, t = 0$.

Proof: Assume that $|Y| > 1$ and $J_{2,s,t} \neq \emptyset$.

Suppose that $J_1 \cup J_{2,s,t}$ is an ideal. Let $\alpha \in J_{2,s,t}$. Then $|X\alpha| = 2$, $|Y\alpha| = s$ and $|X\alpha \setminus Y| = t$. Since $|X\alpha| = 2$ and $1 \leq |Y\alpha| \leq |X\alpha| = 2$, we have $1 \leq s \leq 2$, so $0 \leq |X\alpha \setminus Y| \leq 1$. Thus $0 \leq t \leq 1$. So there are four possible cases: $s = 2$ and $t = 0$; $s = 2$ and $t = 1$; $s = 1 = t$; or $s = 1$ and $t = 0$.

If $s = 2$ and $t = 1$, then $|X\alpha| = 2 = |Y\alpha|$. Since $Y\alpha \subseteq X\alpha$, we obtain that $X\alpha = Y\alpha$ and thus $t = |X\alpha \setminus Y| = |Y\alpha \setminus Y| = 0$ which is a contradiction.

If $s = 2$ and $t = 0$, then $J_1 \cup J_{2,s,t} = J_1 \cup J_{2,2,0}$. Let $\beta \in J_{2,2,0}$. So $|X\beta| = 2 = |Y\beta|$ and $Y\beta \subseteq Y$, thus we can write

$$\beta = \begin{pmatrix} A & B \\ a & b \end{pmatrix}$$

where $A \cap Y \neq \emptyset \neq B \cap Y$; $a, b \in Y$. Since $\emptyset \neq Y \cap X$, there exists $c \in B$ and define $\gamma \in S(X, Y)$ by

$$\gamma = \begin{pmatrix} C & X \setminus C \\ a & c \end{pmatrix}$$

where $Y \subseteq C$. So

$$\gamma\beta = \begin{pmatrix} C & X \setminus C \\ a & b \end{pmatrix} \notin J_1 \cup J_{2,2,0}.$$

Then $J_1 \cup J_{2,2,0}$ is not an ideal is a contradiction.

If $s = 1 = t$, then $J_1 \cup J_{2,s,t} = J_1 \cup J_{2,1,1}$. Let $\lambda \in J_{2,1,1}$. So $|X\lambda| = 2$ and $Y\lambda \subseteq Y$, thus we can write

$$\lambda = \begin{pmatrix} A & X \setminus A \\ u & v \end{pmatrix}$$

where $Y \subseteq A$; $u \in Y$, $v \in X \setminus Y$. Since $|Y| > 1$, there exists $u \neq w \in Y$ and define $\mu \in S(X, Y)$ by

$$\mu = \begin{pmatrix} Y & X \setminus Y \\ u & w \end{pmatrix}$$

So

$$\lambda\mu = \begin{pmatrix} A & X \setminus A \\ u & w \end{pmatrix} \notin J_1 \cup J_{2,1,1}.$$

Thus $J_1 \cup J_{2,1,1}$ is not an ideal which is a contradiction.

Therefore, $s = 1$ and $t = 0$.

Conversely, assume that $s = 1$ and $t = 0$. We show that $J_1 \cup J_{2,1,0} = S(3, 2, 1)$.

Let $\alpha \in J_1 \cup J_{2,1,0}$. Then $\alpha \in J_1$ or $\alpha \in J_{2,1,0}$. If $\alpha \in J_1$, then $|X\alpha| = 1 < 3$, $|Y\alpha| \leq |X\alpha| = 1 < 2$ and $|X\alpha \setminus Y| = 1 - 1 = 0 < 1$. Thus $\alpha \in S(3, 2, 1)$. For the

other containment, let $\alpha \in S(3, 2, 1)$. Then $|X\alpha| \leq 2$, $|Y\alpha| = 1$ and $|X\alpha \setminus Y| = 0$. If $|X\alpha| = 1$, then $\alpha \in J_1$. If $|X\alpha| = 2$, $|Y\alpha| = 1$ and $|X\alpha \setminus Y| = 0$, then $\alpha \in J_{2,1,0}$.

Theorem 3.5. If $|Y| > 1$, then $J_1 \cup J_{2,1,0}$ is the unique minimal ideal of $S(X, Y)$.

Proof: Suppose that $|Y| > 1$. By Lemma 3.4, we have $J_1 \cup J_{2,1,0}$ is an ideal of $S(X, Y)$. To show that $J_1 \cup J_{2,1,0}$ is a minimal ideal of $S(X, Y)$, let J be an ideal of $S(X, Y)$ such that $J_1 \subseteq J \subseteq J_1 \cup J_{2,1,0}$. It is clear that $J_1 \cup J$. By assumption we have there exists $\alpha \in J_{2,1,0}$ but $\alpha \in J$. We prove that $J \subseteq J_1$ by supposing this false. Then there exists $\beta \in J$, but $\beta \in J_1$. Since $\alpha, \beta \in J_{2,1,0}$, we can write

$$\alpha = \begin{pmatrix} A & X \setminus A \\ a & b \end{pmatrix}$$

where $Y \subseteq A$; $a, b \in Y$ and

$$\beta = \begin{pmatrix} B & X \setminus B \\ a' & c \end{pmatrix}$$

where $Y \subseteq B$; $a', c \in Y$. Let $\gamma, \theta \in S(X, Y)$ be defined by

$$\gamma = \begin{pmatrix} A & X \setminus A \\ u & v \end{pmatrix}, \quad \theta = \begin{pmatrix} a & X \setminus \{a\} \\ a & b \end{pmatrix}$$

where $u \in B \cap Y$, $v \in X \setminus B$. so

$$\begin{aligned} \gamma\beta\theta &= \begin{pmatrix} A & X \setminus A \\ u & v \end{pmatrix} \begin{pmatrix} B & X \setminus B \\ a' & c \end{pmatrix} \begin{pmatrix} a & X \setminus \{a\} \\ a & b \end{pmatrix} \\ &= \begin{pmatrix} A & X \setminus A \\ a & b \end{pmatrix} = \alpha. \end{aligned}$$

Then $\alpha = \gamma\beta\theta \in J$, which is a contradiction. Hence, $J_1 \cup J_{2,1,0}$ is a minimal ideal of $S(X, Y)$. Now, we show that $J_1 \cup J_{2,1,0}$ is a unique minima ideal of $S(X, Y)$. Let $M = J_1 \cup J_{2,1,0}$ and N be a minimal ideal of $S(X, Y)$. Since N is an ideal of $S(X, Y)$, we get that $N = K(Z)$ for some $\emptyset \neq Z \subseteq S(X, Y)$. Since $J_1 \cap N$, there exists $\alpha \in N$ with $|X\alpha| \geq 2$. Since $\alpha \in N = K(Z)$, we obtain that $|X\alpha| \leq |X\beta|$, $|Y\alpha| \leq |Y\beta|$ and $|X\alpha \setminus Y| \leq |X\beta \setminus Y|$ for some $\beta \in Z$. Let $\gamma \in J_{2,1,0}$. Then $|X\gamma| = 2$ and so

$|X\gamma| = 2 = |X\alpha| \leq |X\beta|$. Since $|Y| > 1$, we have $|Y\gamma| = 1 \leq |Y\alpha| \leq |Y\beta|$ and $|X\alpha \setminus Y| = 0 \leq |X\alpha \setminus Y| \leq |X\beta \setminus Y|$. Then $\gamma \in K(Z) = N$. Thus $J_{2,1,0} \subseteq N$ and so $J_1 \cup J_{2,1,0} \in N$ which implies that $M = N$.

Lemma 3.6. $J_1 \cup J_2 \cup \dots \cup J_k$ is an ideal of $S(X, Y)$ where $1 \leq k \leq n$.

Proof: Let $\alpha \in J_1 \cup J_2 \cup \dots \cup J_k$ and $\beta, \gamma \in S(X, Y)$. Then $\alpha \in J_{i_0}$ for some $1 \leq i_0 \leq k$ and thus $|X\alpha| = i_0$. Since $X\beta\alpha\gamma = (X\beta)\alpha\gamma \subseteq X\alpha\gamma$, we get that $|X\beta\alpha\gamma| \leq |X\alpha\gamma| \leq |X\alpha| = i_0$. Then $|X\beta\alpha\gamma| = p$ for some $1 \leq p \leq i_0$. Hence $\beta\alpha\gamma \in J_p \subseteq J_1 \cup J_2 \cup \dots \cup J_k$.

An ideal $I \emptyset S(X, Y)$ of $S(X, Y)$ is a maximal ideal if J is an ideal such that $I \subseteq J \emptyset S(X, Y)$, then $I = J$.

Lemma 3.7. Let S be a semigroup with identity 1. If S has a maximal ideal, then it is unique.

Proof: suppose that S has a maximal ideal, say M . Let M' be a maximal ideal of S . It is clear that $M \cup M'$ is an ideal and $1 \notin M \cup M'$. Since $M \subseteq M \cup M'$ and M is a maximal ideal, we have $M \cup M' = M$. Similarly, we have $M \cup M' = M'$. So $M = M \cup M' = M'$ and therefore, S has a unique maximal ideal of S .

If $|X| = |Y| = 1$, then $S(X, Y) = G(X, Y)$. Thus $S(X, Y) \setminus G(X, Y) = \emptyset$. So we consider the case $|X| > 1$.

Theorem 3.8. If $|X| > 1$, then $S(X, Y) \setminus G(X, Y)$ is a unique maximal ideal of $S(X, Y)$.

Proof Let $a \in Y$ and α be the constant map with $X\alpha = \{a\}$. Then $\alpha \in S(X, Y) \setminus G(X, Y)$, so $S(X, Y) \setminus G(X, Y) \neq \emptyset$. By Lemma 3.6, we have $S(X, Y) \setminus G(X, Y) = S(X, Y) \setminus J_n = J_1 \cup J_2 \cup \dots \cup J_{n-1}$ is an ideal of $S(X, Y)$. We show that $S(X, Y) \setminus G(X, Y)$ is a maximal ideal of $S(X, Y)$. Let I be an ideal of $S(X, Y)$ such that $S(X, Y) \subseteq I \emptyset S(X, Y)$. We prove that $I = S(X, Y) \setminus G(X, Y)$ by supposing this is not true. Then there exist $\alpha \in I$ but $\alpha \notin S(X, Y) \setminus G(X, Y)$, i.e., $\alpha \in G(X, Y)$. Since $G(X, Y)$ is a group, we obtain that $\alpha^{-1} \in G(X, Y)$ and $1_X = \alpha\alpha^{-1} \in I$. Thus $I = S(X, Y)$ which is a contradiction. Therefore $I = S(X, Y) \setminus G(X, Y)$. So $S(X, Y) \setminus G(X, Y)$ is a maximal ideal of $S(X, Y)$. By Lemma 3.7, we obtain that

$S(X, Y) \setminus G(X, Y)$ is a unique maximal ideal of $S(X, Y)$.

Let ρ be a congruence on a semigroup S . We recall that ρ is a maximal congruence if δ is a congruence on S with $\rho \emptyset \delta \subseteq S \times S$ implies $\delta = S \times S$.

Theorem 3.9 Let $S = S(X, Y)$ and $G = G(X, Y)$. Then $\rho = (S \setminus G \times S \setminus G) \cup (G \times G)$ is a maximal congruence on S .

Proof It is clear that ρ is a equivalence relation on S . Let $\alpha, \beta, \gamma \in S$ and $(\alpha, \beta) \in \rho$. Then $(\alpha, \beta) \in (S \setminus G \times S \setminus G)$ or $(\alpha, \beta) \in G \times G$. If $(\alpha, \beta) \in (S \setminus G \times S \setminus G)$, then $\gamma\alpha, \alpha\gamma, \gamma\beta, \beta\gamma \in S \setminus G$ since $S \setminus G$ is an ideal of $S(X, Y)$. Thus $(\gamma\alpha, \gamma\beta), (\alpha\gamma, \beta\gamma) \in (S \setminus G) \times (S \setminus G) \subseteq \rho$. If $(\alpha, \beta) \in G \times G$, we consider two cases.

Case 1: $\gamma \in S \setminus G$. Since $S \setminus G$ is an ideal, we have $(\gamma\alpha, \gamma\beta), (\alpha\gamma, \beta\gamma) \in (S \setminus G) \times (S \setminus G) \subseteq \rho$.

Case 2: $\gamma \in G$. Then $\alpha, \beta, \gamma \in G$ and G is a group, so we obtain that $\gamma\alpha, \alpha\gamma \in G$ and $\gamma\beta, \beta\gamma \in G$. Thus $(\gamma\alpha, \gamma\beta), (\alpha\gamma, \beta\gamma) \in G \times G \subseteq \rho$.

Next, we show that ρ is a maximal congruence on S . Let δ be a congruence on S such that $\rho \emptyset \delta \subseteq S \times S$. Since $\rho \emptyset \delta$, there exists $(\alpha, \beta) \in (\delta \setminus \rho)$ with $\alpha \in S \setminus G$ and $\beta \in G$. Let k be the order of β . Then $1_X = \beta^k \delta \alpha^k$ where $\alpha^k \in S \setminus G$ since $S \setminus G$ is an ideal. Now, let $(\lambda, \mu) \in S \times S$. So $\lambda \delta \alpha^k \lambda$ and $\mu \delta \alpha^k \mu$ where $\alpha^k \lambda, \alpha^k \mu \in S \setminus G$ so $\alpha^k \lambda \rho \alpha^k \mu$. Since $\rho \subseteq \delta$, we have $\alpha^k \lambda \delta \alpha^k \mu$. Thus $\lambda \delta \mu$ and $\delta = S \times S$ as required.

ACKNOWLEDGMENT

This research is supported by the Damrongchaitham Foundation. The first author would like to thank the Graduate School Chiang Mai University.

REFERENCES

- [1] W. Choomanee, P. Honyam and J. Sanwong, Regularity in semigroups of transformations with invariant sets, IJPAM., 87 (2013), No. 1, 151- 164.
- [2] A. H. Clifford and G. B. Preston, The algebraic theory of semigroups. Vol. I and II, Mathematical Surveys, No. 7, American Mathematical Society, Providence, R.I. 1961 and 1967.
- [3] P. A. Grillet, Intersections of Maximal Ideals in Semigroups, MAA., 76 (1969), No. 5, 503-509.
- [4] P. Honyam and J. Sanwong, Semigroup of transformations with invariant sets, J. Korean Math. Soc., 48 (2011), No. 2, 289-300.
- [5] K. D. Magill Jr., Subsemigroups of $S(X)$, Math. Japon., 11 (1966), 109-115.
- [6] A. I. Malcev, Symmetric groupoids, Mat. Sbornik N. S. 31(73) (1952), 136-151.
- [7] S. Nenthein, P. Youngkhong and Y. Kemprasit, Regular elements of some transformation semigroups, Pure Math. Appl., 16 (2005), No. 3, 307-314.